

Mini-Review

A temperature-sensitive TRP ion channel, Painless, functions as a noxious heat sensor in fruit flies

Takaaki Sokabe¹ and Makoto Tominaga^{1,2}

¹Section of Cell Signaling; Okazaki Institute for Integrative Bioscience (National Institute for Physiological Sciences); National Institutes of Natural Sciences; Okazaki, Japan;

²Department of Physiological Sciences; The Graduate University for Advanced Studies; Okazaki, Japan

Abbreviations: TRP, transient receptor potential; HEK293 cells, human embryonic kidney 293 cells

Key words: TRP channel, temperature sensation, noxious heat, drosophila, Ca²⁺ ion

Animals must be capable of sensing hazardous temperatures to avoid physical injury. Recent progress has revealed the molecular mechanisms underlying this capability. This essential function requires a subset of the Transient Receptor Potential (TRP) channel family in both mammals and *Drosophila*. We recently showed that a *Drosophila* TRP channel, dubbed Painless, possesses heat sensitivity that is essential for avoidance of noxious heat. The temperature threshold for Painless activation is consistent with the temperatures that cause avoidance behaviors in vivo, indicating that Painless acts as a primary noxious heat detector in *Drosophila*. In this review, we summarize the properties of temperature-sensitive TRP channels, including Painless, in fruit flies.

Temperature-Sensitive TRP Channels are Physiological Thermosensors

The ability to sense environmental temperature is necessary to discriminate between permissive habitats and hazardous environments that cause injury. Recently, it has been revealed that several members of the TRP ion channel super family are activated by temperature changes. The temperature sensitivity of molecules is usually evaluated with their Q₁₀ values, which are defined as ratios of reactions at 10-degree increments. Many enzymes and channels have Q₁₀ values less than 3,¹ whereas the thermosensitive TRP channels (thermoTRPs) have extremely high values greater than 10. A number of reports clearly demonstrate that thermal activation of these TRP channels contributes to various temperature-dependent responses in vivo, such as thermosensation, thermotaxis and regulation of cellular/tissue functions at physiological body temperature. So far, nine TRP channels have been reported to respond to a physiological range of temperatures in mammals.² TRPV1 and TRPV2 expressed

in nociceptive neurons are activated by heat (>43°C and >52°C, respectively), and TRPV1-null mice have defects in sensing noxious heat.³ TRPV3 and TRPV4 are predominantly expressed in skin keratinocytes rather than in sensory neurons, and gene knock-out of each channel causes abnormal thermotaxis in vivo.^{4,5} TRPM8, which senses cold temperatures (<27°C),^{6,7} is expressed in nociceptive and non-nociceptive neurons and its loss impairs cold sensitivity.⁸⁻¹² TRPA1 is expressed in nociceptive neurons¹³ and acts as a sensor for various harmful stimuli, whereas its responsiveness to noxious cold stimuli is controversial even after the analysis of mice lacking the channel.¹⁴⁻¹⁷ Other thermoTRPs, TRPM2, TRPM4 and TRPM5 are not expressed in sensory neurons, and are reportedly involved in several functions at physiological body temperatures including insulin secretion, taste sensation and immune response.¹⁸⁻²⁰ Thus, the thermoTRPs are now considered to be primary elements for temperature-dependent responses.

TRP Channel-Dependent Thermosensation in Fruit Flies

In fruit flies, some TRP channels are important for temperature detection like in mammals. In *Drosophila melanogaster*, thirteen genes encode TRP channels. Five of those channels belong to the TRPA and the TRPC subfamilies, which contribute to temperature-related behaviors (Fig. 1). dTRPA1 was first identified as a heat-activated channel (>27°C) in vitro,²¹ and its knockdown by RNAi²² or gene-knockout²³ revealed the importance of the channel for thermotaxis from warm temperature to a preferred range (18–24°C). dTRPA1 proteins are expressed in anterior cell neurons, two pairs of neurons at the brain's anterior.²⁴ Pyrexia is a noxious heat-sensitive TRPA channel with high potassium permeability, and it may protect flies from high temperature stress.²⁵ Pyrexia-expressing sensory neurons are widely distributed throughout the body. Another TRPA subtype, dubbed Painless, was found to be essential for avoidance of noxious heat above 40°C in *Drosophila* larvae and adults.^{26,27} The protein was found at the tips of the dendritic arbor in a small subset of multidendritic sensory neurons which resemble our peripheral afferent neurons. More recently, TRP and TRPL (belonging to the TRPC subfamily), which are critical for vision in *Drosophila*, were reported to be involved in cold avoidance.²⁸ This evidence strongly demonstrates that certain TRP channels share a common role in

Correspondence to: Takaaki Sokabe; Section of Cell Signaling; Okazaki Institute for Integrative Bioscience; National Institutes of Natural Sciences; Okazaki, Aichi 444-8787 Japan; Tel.: +81.564.59.5287; Fax: +81.564.59.5285; Email: sokabe@nips.ac.jp/ Makoto Tominaga; Section of Cell Signaling; Okazaki Institute for Integrative Bioscience; National Institutes of Natural Sciences; Okazaki, Aichi 444-8787 Japan; Tel.: +81.564.59.5286; Fax: +81.564.59.5285; Email: tominaga@nips.ac.jp

Submitted: 12/19/08; Accepted: 12/23/08

Previously published online as a *Communicative & Integrative Biology* E-publication: <http://www.landesbioscience.com/journals/cib/article/7708>

temperature sensation between distantly related species. Interestingly, in flies, four channels are classified in the TRPA subfamily and three of them are known to contribute to thermosensation. Therefore, the TRPA subfamily may play a central role in temperature detection and related behaviors in insects.

Noxious Heat Sensitivity of Painless

Drosophila melanogaster prefer temperatures in the range of 18–24°C. They quickly escape from noxious temperatures above 40°C. Wild-type larvae display stereotypical ‘rolling behavior’, which is defined as a vigorous rolling movement lateral to the body axis, within one second of being touched with a heated probe.²⁶ Genetic screening to obtain mutants that have abnormal response against noxious heat identified *painless* mutants. These mutant larvae required much longer times to escape from a heated probe, indicating that the *painless* gene is essential for heat avoidance. The mRNA and the protein appeared to be localized in the tips of multidendritic sensory neurons of the peripheral nervous system, which resemble peripheral sensory neurons in mammals. The firing of these neurons increased upon heating over 38°C in wild type neurons, whereas such increases were not seen in the *painless* mutant. The Painless protein is predicted to have ankyrin repeats in the N-terminal region, and six transmembrane domains with one putative pore-forming region between transmembrane domains 5 and 6, all of which are common structures for the TRP ion channel super family. Since two *Drosophila* TRPA channels, dTRPA1 and Pyrexia, have been shown to have temperature sensitivity both in vitro and in vivo, it has been inferred that Painless might also be a heat-sensitive TRP channel. Indeed, we first revealed that the channel actually functions as a thermosensor.²⁹ Painless expressed in human embryonic kidney (HEK) 293 cells showed transient inward currents upon heating at negative membrane potentials, and the temperature thresholds for activation (42–44°C) were consistent with the temperatures that cause avoidance behavior in vivo.²⁶ Painless displayed heat responses in a cell-free excised membrane patch, indicating that Painless may sense heat directly, without utilizing intracellular signaling molecules. These in vitro analyses clearly demonstrate that in *Drosophila*, Painless itself is a primary heat detector, the activation of which raises neural activity against dangerous heat.

Functional Regulation of Painless by Ca²⁺

A unique property of Painless is the extremely high and selective permeability of Ca²⁺, which is almost 42 times higher than that of sodium.²⁹ Intriguingly, the various activation properties of the channel are affected by Ca²⁺, all of which seem to be correlated with the flies’ physiology. This interpretation is supported by the following observations. First, Ca²⁺ enables Painless to be heat-sensitive. Painless failed to respond to heat in the absence of intracellular and extracellular Ca²⁺, whereas 200 nM intracellular Ca²⁺ restored the heat-dependent activation. The mammalian thermoTRPs such as TRPM4, TRPM5 and TRPA1 are activated by intracellular Ca²⁺,^{18,30–32} whereas Painless requires Ca²⁺ as a co-agonist for heat-evoked activation. A similar concept has been reported in TRPM8, in that intracellular Ca²⁺ supports robust icilin-evoked responses.³³ Intracellular Ca²⁺ (100–200 nM) was sufficient for Painless activation, a concentration close to the reported value in the terminal and dorsal organs of the larval head.³⁴ Second, Ca²⁺ accelerates Painless




Figure 1. TRP channels and thermosensation in *Drosophila*. Flies utilize TRP channels to sense environmental temperatures. dTRPA1 is activated above 27°C and contributes to avoidance of warmer temperatures. Pyrexia is activated around 40°C and has a higher potassium permeability. Channel activation prevents paralysis during high temperature stress. Painless is activated above 40°C, and is essential for avoiding hazardous temperatures. These channels belong to the TRPA subfamily and possess temperature sensitivity. TRP and TRPL channels, both of which belong to the TRPC subfamily, are involved in cool temperature avoidance. However, the temperature sensitivity of these channels is unknown. The preferred range of *Drosophila melanogaster* (18–24°C) is indicated by dotted lines.

activation. Channel activation is rapid in the presence (but not in the absence) of physiological levels of intracellular Ca²⁺, which reflects the fly larva’s rapid rolling response when challenged by a heated probe.²⁶ Third, Ca²⁺ sensitizes Painless to heat. The temperature threshold for activation was ~42.6°C in the presence of intracellular Ca²⁺ regardless of the presence of extracellular Ca²⁺, which was significantly lower than that in the presence of extracellular Ca²⁺ alone (~44.1°C). Quick avoidance from a heated probe occurred around 42°C in fly larvae.²⁶ Therefore, the in vivo temperature threshold seems close to that obtained in the presence of intracellular Ca²⁺. Fourth, Ca²⁺ sensitizes Painless upon repetitive heating. Significant reduction in the temperature thresholds by more than 1°C was observed in the presence of extracellular and intracellular Ca²⁺, but not in the presence of intracellular Ca²⁺ alone upon repeated heating, suggesting that extracellular Ca²⁺ and/or Ca²⁺ influx play a role in the sensitization. This is reasonable because flies are able to escape from a hazardous heat source in less time after a second exposure. Sensitization upon repetitive heating is a common feature in TRPV1, TRPV2 and TRPV3,^{35,36} although the underlying mechanism is still unknown, including the requirement for Ca²⁺.

Ca²⁺-dependent regulation is partly ascribed to the properties of the N-terminal region of Painless.²⁹ Specific amino acid substitution (N363A), which is located in the ankyrin repeat domain, displayed small heat-evoked currents, increased temperature thresholds, and higher intracellular Ca²⁺ requirement with a reduced Hill coefficient. These results suggest that N363 is a key residue in Ca²⁺ sensitivity of Painless. Interestingly, the amino acid sequence, which includes N363, has partial similarity to the EF-hand-like domain in mammalian TRPA1. The EF-hand-like motif is reported to be responsible for Ca²⁺-evoked activation of human TRPA1 which was impaired by an amino acid substitution.^{31,32} However, the function of a Ca²⁺-regulatory site in Painless seems different from that of the EF-hand-like motif. TRPA1 requires Ca²⁺ at a micromolar range for its activation, whereas Painless was not activated by intracellular Ca²⁺

alone and requires much less Ca^{2+} for its modulation. Furthermore, mutation of N356, S357 or D366 did not affect the properties of Painless, while the corresponding amino acids in the EF-hand-like motif are critical for Ca^{2+} -dependent TRPA1 activation. Such Ca^{2+} -dependency upon thermal activation is unique to Painless.

Is Painless a Polymodal Receptor?

Our investigation demonstrated that Painless did not respond to typical stimuli known to activate mammalian thermoTRPs.²⁹ This is somewhat surprising because most mammalian thermoTRPs are activated by various stimuli in addition to temperature,³⁷⁻³⁹ and Painless has also been reported to be involved in sensitivity to allyl isothiocyanate, a wasabi ingredient, sugar or mechanical stimuli in vivo.^{26,40,41} Expression of *painless* was seen in a subset of sensory neurons among gustatory bristles located in the labial palpus, the legs, the wings, and the internal pharyngeal sensory structures in adults, all of which have been known to contribute to gustatory detection.⁴⁰ Expression of *painless* was also observed in mechanically-sensitive Johnston's organ and the central nervous system including the mushroom body,²⁶ where Painless could be activated by mechanical stimuli or endogenous ligands, respectively. Therefore, Painless possibly senses multiple stimuli other than heat, which might require endogenous accessory proteins, heteromeric channel formation, splice variants and/or phosphorylation. GPCR signaling might also regulate Painless function since the G_q -PLC pathway has been implicated in modulation of dTRPA1-dependent thermosensation.²³ These observations must be confirmed under more physiological conditions. It should be noted that heat activation of Painless was inhibited by ruthenium red and camphor,²⁹ which was reminiscent of the properties of mammalian TRPA1. This feature suggests some common mechanisms for channel activation within mammalian and *Drosophila* TRPA channels.

Thus far, no chemical or other physical stimuli have been available for activation of invertebrate thermoTRPs. Identification of new activators would be useful to control insects' behavior as the TRP channels primarily contribute to sensing mechanisms in invertebrates as well as in vertebrates. Camphor, a wood derivative from camphor laurel, has been used as a repellent for pests and proved to be effective for mosquitoes.⁴² It would, therefore be intriguing to test the effects of camphor at the behavioral level, in spite of its inhibitory effect on Painless. Since activation of Painless evokes 'avoidance' in larvae and adult flies,^{26,27} this channel is a good target for the development of new insect repellents.

Future Tasks: Analyzing the Mechanism of Thermosensation

Identification of the thermoTRPs sheds light on molecular mechanisms for temperature sensation. However, compared to the recent remarkable progress in mammalian research fields, understanding of the molecular properties and the physiological roles of thermoTRPs in ectotherms and invertebrates is quite limited, although the temperature sensation must be more important for them than for endotherms. Recent studies, including our own, provide clear evidence that fruit flies utilize specific channels to detect warm to hot temperatures, which emphasizes the significance of the evolutionarily conserved property of the TRP channels. In contrast, little is known about the mechanisms for cold sensation in *Drosophila*. They avoid undesirable cold temperatures ($<18^\circ\text{C}$) and some specific neurons

appear to respond to cold exposure.³⁴ Therefore, *Drosophila* must have specific cold sensor(s) like TRPM8 in mammals. In addition to TRP and TRPL channels,²⁸ it has been reported that histamine signaling molecules including a histamine-gated chloride channel, and a cyclic AMP-PKA pathway are involved in temperature preference and cold tolerance in *Drosophila*.^{43,44} Furthermore, a recent report demonstrates that TRPA1 modulation via a G_q -PLC pathway participates in discriminating cooler temperatures in an optimal range ($18\text{--}24^\circ\text{C}$).²³

We have clarified the activation properties of Painless, and they are better understood than any other insect channel. Given that TRP channels play a primary role in sensation, elucidation of the channel properties in various insects would be beneficial to our understanding and facilitate control of their physiological responses. Genetic and behavioral analyses could reveal the basic mechanisms for temperature sensation in fruit flies, which may help us understand the complicated thermoregulatory system in mammals. Comparison of the function of the thermoTRPs among various species may also help clarify how specific channels are activated by temperature changes.

Acknowledgements

This work was supported by grants from the Japan Society for the Promotion of Science in Japan (Takaaki Sokabe) and from the Ministry of Education, Culture, Sports, Science and Technology in Japan (Makoto Tominaga).

References

- Hille B. Ion Channels of Excitable Membrane, 3rd Ed. Sunderland: Sinauer Associates, Inc 2001.
- Bandell M, Macpherson LJ, Patapoutian A. From chills to chilis: Mechanisms for thermosensation and chemesthesis via thermoTRPs. *Curr Opin Neurobiol* 2007; 17:490-7.
- Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeit KR, et al. Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 2000; 288:306-13.
- Moqrich A, Hwang SW, Earley TJ, Petrus MJ, Murray AN, Spencer KS, et al. Impaired thermosensation in mice lacking TRPV3, a heat and camphor sensor in the skin. *Science* 2005; 307:1468-72.
- Lee H, Iida T, Mizuno A, Suzuki M, Caterina MJ. Altered thermal selection behavior in mice lacking transient receptor potential vanilloid 4. *J Neurosci* 2005; 25:1304-10.
- McKemy DD, Neuhauser WM, Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* 2002; 416:52-8.
- Peier AM, Moqrich A, Hergarden AC, Reeve AJ, Andersson DA, Story GM, et al. A TRP channel that senses cold stimuli and menthol. *Cell* 2002; 108:705-15.
- Bautista DM, Siemens J, Glazer JM, Tsuruda PR, Basbaum AI, Stucky CL, et al. The menthol receptor TRPM8 is the principal detector of environmental cold. *Nature* 2007; 448:204-8.
- Colburn RW, Lubin ML, Stone DJ Jr, Wang Y, Lawrence D, D'Andrea MR, et al. Attenuated cold sensitivity in TRPM8 null mice. *Neuron* 2007; 54:379-86.
- Dhaka A, Murray AN, Mathur J, Earley TJ, Petrus MJ, Patapoutian A. TRPM8 is required for cold sensation in mice. *Neuron* 2007; 54:371-8.
- Takashima Y, Daniels RL, Knowlton W, Teng J, Liman ER, McKemy DD. Diversity in the neural circuitry of cold sensing revealed by genetic axonal labeling of transient receptor potential melastatin 8 neurons. *J Neurosci* 2007; 27:14147-57.
- Dhaka A, Earley TJ, Watson J, Patapoutian A. Visualizing cold spots: TRPM8-expressing sensory neurons and their projections. *J Neurosci* 2008; 28:566-75.
- Story GM, Peier AM, Reeve AJ, Eid SR, Mosbacher J, Hricik TR, et al. ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures. *Cell* 2003; 112:819-29.
- Kwan KY, Allchorne AJ, Vollrath MA, Christensen AP, Zhang DS, Woolf CJ, et al. TRPA1 contributes to cold, mechanical and chemical nociception but is not essential for hair-cell transduction. *Neuron* 2006; 50:277-89.
- Bautista DM, Jordt SE, Nikai T, Tsuruda PR, Read AJ, Poblete J, et al. TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell* 2006; 124:1269-82.
- Jordt SE, Bautista DM, Chuang HH, McKemy DD, Zygmunt PM, Hogestatt ED, et al. Mustard oils and cannabinoids excite sensory nerve fibres through the TRP channel ANKTM1. *Nature* 2004; 427:260-5.
- Sawada Y, Hosokawa H, Hori A, Matsumura K, Kobayashi S. Cold sensitivity of recombinant TRPA1 channels. *Brain Res* 2007; 1160:39-46.

18. Talavera K, Yasumatsu K, Voets T, Droogmans G, Shigemura N, Ninomiya Y, et al. Heat activation of TRPM5 underlies thermal sensitivity of sweet taste. *Nature* 2005; 438:1022-5.
19. Togashi K, Hara Y, Tominaga T, Higashi T, Konishi Y, Mori Y, et al. TRPM2 activation by cyclic ADP-ribose at body temperature is involved in insulin secretion. *EMBO J* 2006; 25:1804-15.
20. Cheng H, Beck A, Launay P, Gross SA, Stokes AJ, Kinet JP, et al. TRPM4 controls insulin secretion in pancreatic beta-cells. *Cell Calcium* 2007; 41:51-61.
21. Viswanath V, Story GM, Peier AM, Petrus MJ, Lee VM, Hwang SW, et al. Opposite thermosensor in fruitfly and mouse. *Nature* 2003; 423:822-3.
22. Rosenzweig M, Brennan KM, Tayler TD, Phelps PO, Patapoutian A, Garrity PA. The *Drosophila* ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes Dev* 2005; 19:419-24.
23. Kwon Y, Shim HS, Wang X, Montell C. Control of thermotactic behavior via coupling of a TRP channel to a phospholipase C signaling cascade. *Nat Neurosci* 2008; 11:871-3.
24. Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, et al. An internal thermal sensor controlling temperature preference in *Drosophila*. *Nature* 2008; 454:217-20.
25. Lee Y, Lee J, Bang S, Hyun S, Kang J, Hong ST, et al. Pyrexia is a new thermal transient receptor potential channel endowing tolerance to high temperatures in *Drosophila melanogaster*. *Nat Genet* 2005; 37:305-10.
26. Tracey WD Jr, Wilson RI, Laurent G, Benzer S. painless, a *Drosophila* gene essential for nociception. *Cell* 2003; 113:261-73.
27. Xu SY, Cang CL, Liu XF, Peng YQ, Ye YZ, Zhao ZQ, et al. Thermal nociception in adult *Drosophila*: behavioral characterization and the role of the painless gene. *Genes Brain Behav* 2006; 5:602-13.
28. Rosenzweig M, Kang K, Garrity PA. Distinct TRP channels are required for warm and cool avoidance in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 2008; 105:14668-73.
29. Sokabe T, Tsujiuchi S, Kadowaki T, Tominaga M. *Drosophila* painless is a Ca^{2+} -requiring channel activated by noxious heat. *J Neurosci* 2008; 28:9929-38.
30. Nilius B, Mahieu F, Prenen J, Janssens A, Owsianik G, Vennekens R, et al. The Ca^{2+} -activated cation channel TRPM4 is regulated by phosphatidylinositol 4,5-bisphosphate. *EMBO J* 2006; 25:467-78.
31. Zurborg S, Yurgionas B, Jira JA, Caspani O, Heppenstall PA. Direct activation of the ion channel TRPA1 by Ca^{2+} . *Nat Neurosci* 2007; 10:277-9.
32. Doerner JF, Gisselmann G, Hatt H, Wetzel CH. Transient receptor potential channel A1 is directly gated by calcium ions. *J Biol Chem* 2007; 282:13180-9.
33. Chuang HH, Neuhauser WM, Julius D. The super-cooling agent icilin reveals a mechanism of coincidence detection by a temperature-sensitive TRP channel. *Neuron* 2004; 43:859-69.
34. Liu L, Yermolaieva O, Johnson WA, Abboud FM, Welsh MJ. Identification and function of thermosensory neurons in *Drosophila* larvae. *Nat Neurosci* 2003; 6:267-73.
35. Xu H, Ramsey IS, Kotecha SA, Moran MM, Chong JA, Lawson D, et al. TRPV3 is a calcium-permeable temperature-sensitive cation channel. *Nature* 2002; 418:181-6.
36. Caterina MJ, Rosen TA, Tominaga M, Brake AJ, Julius D. A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature* 1999; 398:436-41.
37. Ramsey IS, Delling M, Clapham DE. An introduction to TRP channels. *Annu Rev Physiol* 2006; 68:619-47.
38. Tominaga M. The Role of TRP Channels in Thermosensation. In: TRP ion channel function in sensory transduction and cellular signaling cascades (Liedtke WB, ed). New York: CRC Press 2007; 271-86.
39. Dhaka A, Viswanath V, Patapoutian A. TRP ion channels and temperature sensation. *Annu Rev Neurosci* 2006.
40. Al-Anzi B, Tracey WD Jr, Benzer S. Response of *Drosophila* to wasabi is mediated by painless, the fly homolog of mammalian TRPA1/ANKTM1. *Curr Biol* 2006; 16:1034-40.
41. Xu J, Sornborger AT, Lee JK, Shen P. *Drosophila* TRPA channel modulates sugar-stimulated neural excitation, avoidance and social response. *Nat Neurosci* 2008; 11:676-82.
42. Gillij YG, Gleiser RM, Zygadlo JA. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. *Bioresour Technol* 2007.
43. Hong ST, Bang S, Paik D, Kang J, Hwang S, Jeon K, et al. Histamine and its receptors modulate temperature-preference behaviors in *Drosophila*. *J Neurosci* 2006; 26:7245-56.
44. Hong ST, Bang S, Hyun S, Kang J, Jeong K, Paik D, et al. cAMP signalling in mushroom bodies modulates temperature preference behaviour in *Drosophila*. *Nature* 2008; 454:771-5.